

MACRO-AGGREGATE STABILITY OF SOILS UNDER SELECTED PARENT MATERIALS IN IMO STATE, NIGERIA.

Ihem, E.E., Nkwopara, U.N., Ndukwu, B.N., Onwudike, S.U.

Department of Soil Science and Technology, Federal University of Technology, P.M.B 1526, Owerri, Imo State.

*Correspondence: eeihem2000@yahoo.com

Abstract

We investigated the effect of macro-aggregate stability of soils formed over selected parent materials at three locations in Imo State: Ihiagwa, Egbema and Okigwe representing the three parent materials (Coastal Plain Sand, Alluvium false-bedded sandstone respectively). Target soil survey sampling technique was used in locating pedons. Two pedon were dug in each of the locations and soil samples were taken for routine analysis on some soil physico-chemical properties and aggregate stability. Soil data were subjected to mean and analysis of variance (ANOVA). Means were separated using least significant difference at 5% and 1% levels of probability. Two indices were used to determine the macro-aggregate stability: water stable aggregate (WSA) and mean weight diameter (MWD). Results showed significant differences ($p \leq 0.05$) on the effect of parent materials on soil physico-chemical properties. While $WSA > 0.3\text{mm}$ positively correlated with effective cation exchange capacity (ECEC) ($r=0.789$), at ($P=0.01$) and total exchange basis (TEB) ($r=0.720$, at $P=0.01$). MWD correlated positively with clay ($r=0.504$, at $P=0.05$) and negatively with pH ($r=0.56$, at $P=0.05$).

Keywords: Aggregate stability, pedon, parent materials, mean weight diameter.

Introduction

Ability of soil aggregates to resist disruption when outside forces (usually associated with water) are applied, (NCRS, 1999) is a very important attribute in erosion-prone landscapes.

Soil aggregate stability influences a wide range of physical and biochemical processes of agricultural soils especially in the tropics. It is a crucial and valuable soil property affecting soil sustainability and crop production (Amezketta, 1999).

The physical and chemical properties of individual aggregate size fractions differs from one another. The contents of clay, organic matter and exchangeable cations for example, often differ in one aggregate size fraction than those in another (Cambardella and Elliot, 1993; Cruvinelet *et al.*, 1999). The stability of macro aggregates varies with the changes in organic matter content or in management practices (Chenuet *et al.*, 2000; Boix-fayoset *et al.*, 2001; Six *et al.*, 2004; Noellemeyeret *et al.*, 2008). This is because macro aggregates are stabilized by such transient or temporary binding agents as roots; hyphae and microbes as well as plant-dried polysaccharides.

Consequently, the stability of the soil aggregation is very reliant on the type of minerals that are present in the soil. Parent materials have a great influence on soil formation, through their mineralogical composition, texture and their stratification as well as other soil properties (Brady and Weil, 1999; Akamigbo, 2001; Esu, 2005). Certain clay minerals form very stable aggregates while others form weak aggregates that fall apart very easily.

Tropical soils are highly weathered, erodible and instable. Most of the soils are therefore poorly structured.

Previous research works by (Osuji and Onwrenmadu, 2005), (Opara, 2009) were mostly on aggregation of Mediterranean soil and the use of aggregates size as well as enhancing the water stable aggregate of tropical soils were the main focus of their studies but not much has been done in aggregate stability based on the different parent materials.

Therefore, this study investigated the macro-aggregate stability of three soils formed over different parent materials in Imo State.

Specific objectives include to:

- determine the physic-chemical properties of the soils in relation to aggregate stability.
- compare the stability of the different aggregates sizes of the soils.
- determine which aggregate size fraction of the soil that would provide more stability.

Materials and Methods

Study Area

This was carried out in the three locations of Imo State, Nigeria: Ihiagwa (Owerri) lies on latitude $5^{\circ}24'N$ and longitude $7^{\circ}1'E$; Egbema (Ohaji/Egbema) on latitude $5^{\circ}54'N$ and longitude $6^{\circ}72'E$ and Okigwe which lies on latitude $5^{\circ}87'N$ and longitude $7^{\circ}30'E$. The main geological material of Ihiagwa is coastal plain sands Benin formation (Orajaka, 1975); Egbema is of Alluvium and Okigwe soil is formed from false bedded sand stone. The study area lies within the lowland area of Eastern Nigeria (Ofomata, 1975). It is of a humid tropical climate characterized by a bimodal rainfall peak ranging from 2000 – 2500mm annually. Temperatures are usually high and ranges from $26-30^{\circ}C$. The area is dominated by rainforest vegetation whose density has been altered by anthropogenic activities such as oil activities, automobile related

serving centres, deforestation, farming etc. The socio-economic activities of the area include hunting, fishing, farming, oil business etc.

Field Studies

Two profile pits were dug at each selected site to give a total of six (6) profile pits from the three different geological material. Coastal plain sands (Ihiagwa), Imo Clay-Shale (Egbema) and False bedded sand stone (Okigwe). Profile pits were dug, examined following the guideline according to FAO (2006). Sampling was done based on horizon differentiation.

Soil samples were air-dried, crushed and sieved using 2-mm sieve, preparatory for laboratory analysis and the remaining samples for macroaggregate stability determination.

Particle Size distribution was determined by hydrometer method according to the procedure of Gee and Or (2002)

Moisture Content was determined gravimetrically by Cater method (1993) thus:

$$\theta MS = \frac{\text{Wet Soil Sample} - \text{Dry Soil sample}}{\text{Dry Soil Sample}} \times \frac{100}{1}$$

Where θMS gravimetric moisture content (saturated)%.

Bulk Density was determined by using core sample method (Grossman and Reinsch, 2002). **Soil pH** was measures in 1:2.5 soil-water suspension methods using a standard pH meter (Thomas, 1996). **Organic carbon** was determined by Walkley and Black wet digestion method (Nelson and Sommers, 1982), while the organic matter was obtained by multiplying the result from organic carbon with the Van Bemmele factor of 1.724. Exchangeable bases were estimated by the neutral ammonium acetate procedure (Thomas, 1982).

Exchangeable acidity (H^+ and Al^{3+}) were determined using titrimetric method (McLean, 1982). Total nitrogen was determined using microkjedahl digestion method (Bremner, 1996).

Available phosphorus was estimated according to the method of Olson and Somner (1990). Macro-aggregate stability was determined using modified methods by Kemper and Rosenau (1986) and NCRS (1996). Wet and dry sieving methods were carried out using a nest of five sieves with aperture sizes of: 4.75, 2.0, 1.1, 0.85 and 0.3mm. Two indices were used to express the aggregate stability;

Mean weight diameter: $MWD = \sum_{i=2}^n \bar{x}_i W_i$

Where \bar{X}_i = the mean diameter of aggregate size fraction i ($i = 1$ to n)

W_i = the proportion of aggregate size fraction 1 to the total sample weight.

Water stable aggregate (WSA) >0.3mm

$$WSA (>0.3mm) = \frac{\text{Weight of dried aggregate-sand}}{\text{weight of Sample-Sand}} \times \frac{100}{1}$$

Higher values of the WSA >0.3mm and MWD indices indicate greater stability.

Data Analysis

Soil data generated were subjected to statistical analysis using analysis of variance (ANOVA), Regression analysis and correlation analysis to estimate the relationship between different aggregate sizes and some soil properties. The LSD method of mean separation was also used to determine the levels of significance of the parameters at P(0.01 and 0.05) with the Gen State Statistical package (Baye *et al.*, 2004).

Results and Discussion

Table 1 shows the physical properties of the studies soils. The sand content was highest in Ihiagwa with a mean value of 909.6 g kg⁻¹ and lower in Egbema (639.0 g kg⁻¹) and least in Okigwe with a mean value of 629.0 g kg⁻¹. The sandiness of the soil is attributes to the sandy nature of the parent material, being derived from coastal plain sand which according to Ogban *et al.* (2001) are characterized by the dominance of sandy textured fragment.

The lowest value of clay content was found in Ihiagwa soils with a mean value of 63.2 g kg⁻¹, while the highest value was found in Okigwe with a mean value of 283.0 g kg⁻¹ and Egbema 238. g kg⁻¹.

The silt content was highest in Egbema with a mean value of 123. 0 g kg⁻¹ and was lowest in Okigwe soils with the mean values of 8.8 g kg⁻¹ and Ihiagwa, 27.2 g kg⁻¹ respectively.

The low clay and silt contents were as a result of high rainfall status of the area which favoured high rate of leaching and the influence of pedogenic process of translocation in which clay and silt contents were moved down the profile horizon (Onweremadu *et al.*, 2007). Akamigbo (1999) also opined that climate factors such as high precipitation and temperature in the soils of southeastern Nigeria could lead to low clay and silt contents of the soils.

The moisture content of all the soils were slightly different with the highest mean values recorded in Okigwe and Ihiagwa (182.9g kg⁻¹) and Egbema recorded the lowest mean value of 123.9 kg.

This could be due to their textural differences which is in line with NCRS (1996) who states that a soil's ability to hold water is affected by soil texture, presence and abundance of rock fragments, soil depth and restrictive layers, also organic matter increases a soil's ability to hold water both directly and indirectly. Meanwhile, the result in table 1 showed that water moisture content increased with depth. This could be as a result of increase in clay content down the soil layers and clay attracts and absorbs water (Calero *et al.*, 2008; Dikinya *et al.*, 2006).

Bulk density values were highest in Egbema soils with a mean value of 1.54g/cm³ and low in both Okigwe and Ihiagwa with a mean value of 1.1g/cm³. The high value of bulk density in Egbema soils could be as a result of the geological formation of Imo Clay Shale which is hard, cracks easily and does

not permit root penetration, while the low bulk densities of Okigwe and Ihiagwa indicated that the soils are not probably compacted and have more porosity (Attah, 2010). This is beneficial to root activity, water infiltration and overall growth of crops. Higher bulk density values could as well be attributed to the activities of oil pollution in the area, such as oil spillage which might cause compaction of soil pores and clogging effects. Seasonal flooding is common in the area, and weight of water impact on these soils results in soil compaction.

Results of the chemical properties of the studied soils are shown in Table 2. The result of the study indicated that soils were generally acidic with Okigwe and Egbema having mean values of 3.85 and 3.83 respectively while Ihiagwa recorded a mean of 5.12. According to Landon (1991), soil pH range of 3.5 is ultra acidic while 5.1 – 5.5 is strongly acidic. The low pH could be as a result of high rainfall.

Table 1: Average values of Physical Properties of the Locations

Location	Horizons	Sand g/kg	Silt g/kg	Clay g/kg	TC	BD g/cm ³	TP %	MC g/kg
Okigwe	AP1	774	98	128	SL	1.16	37.5	128.2
Okigwe	Bt1	654	98	248	SCL	1.16	54.68	174
Okigwe	Bt 2	554	98	348	SCL	1	60.94	207.2
Okigwe	Bt 3	534	58	408	SCL	1.12	56.25	222.2
	Mean	629	8.8	283		1.11	52.3425	182.9
Egbema	AP1	774	138	88	SL	1.78	30.47	86.1
Egbema	Bt1	674	118	208	SCL	1.74	32.19	102.2
Egbema	Bt 2	554	138	308	SCL	1.38	46.09	124.5
Egbema	Bt 3	554	98	348	SCL	1.25	48.83	182.9
	Mean	639.00	123.0	238.0		1.54	39.40	123.9
Ihiagwa	AP1	949.60	27.2	23.2	S	1.16	37.50	128.2
Ihiagwa	Bt1	889.60	47.2	63.2	S	1.16	54.68	174.0
Ihiagwa	Bt 2	909.60	27.2	63.2	S	1.00	60.94	207.2
Ihiagwa	Bt 3	889.60	7.2	103.2	S	1.12	56.25	222.2
	Mean	909.60	27.2	63.2		1.11	52.34	182.9
	LSD(0.05)	89.5	1.73	9.03		0.224	7.3	2.175

TC = textural class, BD = bulk density, TP = total porosity MC = moisture content
LSD 0.05

Table 2: Average Values of Chemical Properties of the Location

Location	Horizon	pH(H ₂ O)	OM	TN	Avail P Mg/kg	Ca	Mg	K	Na	Al Cmo/kg	H	TEB	TEA	ECEC	BS
Okigwe	AP1	5.5	1.33	0.108	37.7	2.4	1.8	0.115	0.246	0	0.72	4.561	0.72	5.28	86.38
Okigwe	Bt1	5	1.02	0.089	38.5	2.4	1.6	0.103	0.233	0.06	0.9	4.336	0.96	5.29	81.92
Okigwe	Bt 2	4.8	0.83	0.0073	35.3	2	1.6	0.096	0.272	0.12	0.92	3.968	0.04	5.01	79.2
Okigwe	Bt 3	4.4	0.76	0.06	33.4	2	0.8	0.086	0.272	0.28	0.28	3.058	1.28	4.44	71.13
	Mean	4.925	0.985	0.0825	36.225	2.2	1.45	0.1	0.25558	0.115	0.705	3.9808	1	5.005	79.69
Egbema	AP1	5.5	1.29	2.22	40.5	2.2	1.6	0.109	0.118	0	0.84	4.027	0.84	4.87	82.75
Egbema	Bt1	5.1	1.14	1.97	40	2	1.4	0.105	0.127	0	0.94	3.632	0.94	4.57	79.47
Egbema	Bt 2	4.8	0.8	1.38	39.3	1.6	0.8	0.093	0.129	0.04	0.96	2.622	1.	3.62	72.43
Egbema	Bt 3	4	0.68	1.17	38.5	1.6	0.8	0.084	0.171	0.13	1.01	2.655	1.14	3.79	70.1
	Mean	4.80	0.98	1.69	39.58	1.85	1.15	0.10	0.14	0.04	0.94	3.23	0.98	4.21	76.19
Ihiagwa	AP1	5.42	0.72	1.24	2.52	1.68	0.26	0.02	0.04	1.44	0.24	2.01	1.68	3.68	54.35
Ihiagwa	Bt1	5.82	0.68	1.17	1.05	2.32	1.06	0.01	0.04	0.88	0.44	3.43	1.32	4.76	72.25
Ihiagwa	Bt 2	2.85	0.52	0.89	1.68	1.68	1.02	0.00	0.00	0.00	0.96	2.73	0.96	3.66	73.77
Ihiagwa	Bt 3	5.53	0.34	0.59	1.92	1.92	3.19	0.00	0.00	0.60	0.92	5.12	1.52	6.65	77.13
	Mean	5.66	0.57	0.97	1.90	1.90	1.38	0.02	0.02	0.73	0.64	3.32	1.37	4.69	69.38
	LSD (0.05)		0.146	0.249	0.359	0.395	1.66	0.044	0.044	0.65	0.489	1.916	0.406	1.75	15.83

OM = organic matter, TN = total nitrogen, Avail. P = available, Ca = calcium, Mg = Magnesium, K= Potassium, Na = sodium, Al = aluminum, H = hydrogen, TEB = total exchangeable bases, TEA = total exchangeable acidity, ECEC = effective cation exchange capacity, BS = bases

LSD 0.05

Table 3: Average Values of Water Aggregates

Water stable aggregate under different (mm) sieve sizes %							
Location	Horizon	Aggreg. 4.75	Aggreg. 2.0	Aggreg. 1.1	Aggreg. 0.825	Aggreg. 0.3	WSA>0.3
Okigwe	AP	1.045	3.893	0.121	1.949	13.16	7.008
Okigwe	Bt ₁	1.046	0.621	0.11	0.804	24.89	2.581
Okigwe	Bt ₂	4.626	1.736	0.121	1.708	48.56	8.191
Okigwe	Bt ₃	0.014	1.142	0.11	3.269	30.38	4.535
	Mean	1.683	1.848	0.116	1.933	29.248	5.579
Egbema	AP	-	0.66	0.205	0.399	23.76	0.664
Egbema	Bt ₁	-	0.22	0.814	2.369	11.76	3.603
Egbema	Bt ₂	-	0.05	0.162	0.723	19.77	1.385
Egbema	Bt ₃	-	0.03	0.665	0.77	21.22	1.465
	Mean	-	0.090	0.462	1.065	19.133	1.779
Ihiagwa	AP	-	2.04	0.161	0.327	13.43	0.528
Ihiagwa	Bt ₁	-	0.007	0.131	0.266	13	0.467
Ihiagwa	Bt ₂	-	0.2	0.473	0.572	26.22	1.245
Ihiagwa	Bt ₃	-	0.191	0.223	0.318	11.87	12.602
	Mean	-	0.625	0.247	0.371	16.130	3.711
	LSD(0.05)	-	1.317	0.404	1.577	14.62	7.11

Table 4: Average Values of Dry Aggregates of the Studied Soils

Dry stable aggregate under different (mm) sieve sizes %							
Location	Horizon	Aggreg. 4.75	Aggreg. 2.0	Aggreg. 1.1	Aggreg. 0.825	Aggreg. 0.3	WSA>0.3
Okigwe	AP1	25.46	17.26	5.83	5.96	30.57	54.51
Okigwe	Bt1	44.03	27.94	12.05	5.12	7.13	89.14
Okigwe	Bt 2	41.52	26.71	9.98	4.94	9.83	83.16
Okigwe	Bt 3	45.63	21.22	13.04	3.07	13.04	85.5
	Mean	39.160	23.160	10.225	4.773	15.143	78.078
Egbema	AP1	0	0.56	3	10.17	63.67	13.73
Egbema	Bt1	0	0.16	2.49	9.29	62.26	11.94
Egbema	Bt 2	0	0.15	2.71	98.5	61.43	11.36
Egbema	Bt 3	0	0.42	3.83	11.25	61.21	21.5
	Mean	0.000	0.323	3.008	9.803	62.143	14.633
Ihiagwa	AP1	1.77	12.48	8.75	6.9	44.49	29.9
Ihiagwa	Bt1	21.86	21.52	8.9	7.74	29.49	60.02
Ihiagwa	Bt 2	26.22	17.32	6.79	5.23	24.3	55.56
Ihiagwa	Bt 3	17.81	15.58	8.32	6.59	30.46	48.28
	Mean	16.915	16.725	8.190	6.615	32.185	48.440
	LSD(0.05)	10.65	4.896	3.327	2.039	9.14	15.96

Table 5: Averages Values of Wet Mean Weight Diameter Aggregates of the Studied Soils

Location	Horizon	Wet MWD aggregate under different (mm) sieve sizes %					MWD
		Aggreg. 4.75	Aggreg. 4.75-2.0	Aggreg. 2.0-1.1	Aggreg. 1.1-0.825	Aggreg. 0.825-0.3	
Okigwe	AP1	0.015	0.009	0.015	0.005	0.003	0.047
Okigwe	Bt1	0.014	0.061	.018	0.018	0.109	0.21
Okigwe	Bt 2	0.033	0.012	0.006	0.016	0.078	0.145
Okigwe	Bt 3	0.779	0.16	0.009	0.015	0.031	0.84
	Mean	0.210	0.036	0.010	0.014	0.055	0.311
Egbema	AP1	-	0.005	0.038	0.096	0.326	0.467
Egbema	Bt1	-	0.01	0.038	0.115	0.286	0.402
Egbema	Bt 2	-	0.001	0.026	10.077	0.366	0.473
Egbema	Bt 3	-	0.007	0.046	0.096	0.299	0.443
	Mean	-	0.006	0.037	0.096	0.319	0.446
Ihiagwa	AP1	-	0.009	0.018	0.023	0.208	0.258
Ihiagwa	Bt1	-	0.008	0.018	0.026	0.214	0.266
Ihiagwa	Bt 2	-	0.007	0.017	0.025	0.245	0.244
Ihiagwa	Bt 3	-	0.019	0.026	0.028	0.166	0.239
	Mean	-	0.011	0.020	0.026	0.252	0.252
	LSD	0.379	0.026	0.0078	0.0158	0.368	0.368
	(0.05)						

Table 6: Averages Values of Dry Mean Weight Diameter Aggregates of the Studied Soils

Location	Horizon	Dry MWD aggregate under different (mm) sieve sizes %					MWDw	PSDI
		Aggreg. 4.75	Aggreg. 4.75-2.0	Aggreg. 2.0-1.1	Aggreg. 1.1-0.825	Aggreg. 0.825-0.3		
Okigwe	AP1	0.6	0.583	0.092	0.057	0.077	1.409	0.967
Okigwe	Bt1	1.039	0.944	0.407	0.049	0.015	2.454	0.914
Okigwe	Bt 2	0.994	0.903	0.115	0.047	0.055	2.139	0.932
Okigwe	Bt 3	0.903	0.717	0.241	0.029	0.005	2.069	0.594
	Mean	0.872	0.787	0.224	0.046	0.038	2.018	0.852
Egbema	AP1	-	0.019	0.047	0.098	0.357	0.521	0.104
Egbema	Bt1	-	0.005	0.039	0.089	0.349	0.482	0.068
Egbema	Bt 2	-	0.005	0.042	0.082	0.344	0.472	0.001
Egbema	Bt 3	-	0.014	0.059	0.108	0.343	0.524	0.448

	Mean	-	0.011	0.047	0.094	0.348	0.500	0.155
Ihiagwa	AP1	-	0.059	0.193	0.066	0.427	0.745	0.999
Ihiagwa	Bt1	-	0.739	0.336	0.074	0.283	0.129	0.764
Ihiagwa	Bt 2	-	0.886	0.266	0.05	0.136	0.34	0.764
Ihiagwa	Bt 3	-	0.602	0.241	0.063	0.173	0.079	0.778
	Mean	-	0.572	0.259	0.063	0.255	0.073	0.826
	LSD(0.05)	0.379	0.327	0.125	0.0196	0.123	0.425	0.333

Table 7: Correlation Matric of Soil Physicochemical Properties

	DSA>0.3	MWDw	PSDI	WSA>0.3	BD	BS	Clay	ECEC	MC	Mg	OM	pH(H ₂ O)	Sand	TEA	TEB	TP	TN
DSA>0.3	1																
MWDw	-0.15	1															
PSDI	0.72**	-0.58*	1														
WSA>0.3	0.34	-0.28	0.33	1													
BD	0.72**	0.32	-0.82**	-0.31	1												
BS	0.18	-0.25	-0.14	0.39	0.25	1											
Clay	0.15	0.50*	-0.24	0.14	-0.07	0.12	1										
ECEC	0.36	-0.34	0.29	0.79**	-0.07	0.58*	-0.15	1									
MC	0.70**	0.05	0.52*	0.45	-0.80**	-0.11	0.31	0.23	1								
Mg	0.17	0.37	0.13	0.81**	0.00	0.65*	0.15	0.93*	0.21	1							
OM	-0.16	-0.07	-0.26	-0.19	0.59*	0.55*	0.04	0.04	-0.74**	-0.04	1						
pH(H ₂ O)	0.02	-0.56	0.31	0.03	-0.08	0.09	-0.90**	0.26	-0.14	0.28	-0.05	1					
Sand	-0.11	-0.47	0.39	-0.04	-0.12	-0.24	-0.97**	0.16	-0.08	0.17	-0.24	0.88	1				
TEA	0.06	0.14	0.33	0.11	0.35	-0.79**	-0.19	0.03	0.37	-0.71**	0.06	0.37	1				
TEB	0.31	-0.39	0.18	0.72**	0.05	0.80**	-0.30	0.95*	0.10	-0.25	0.25	0.24	0.06	-0.29	1		
TP	0.72**	-0.05	0.47	0.29	-0.81**	-0.06	0.30	0.13	0.93**	-0.70	-0.70	-0.05	-0.11	0.24	0.05	1	
TN	0.13	-0.13	0.10	-0.30	-0.03	-0.02	-0.31	0.05	-0.04	-0.02	-0.02	0.40	0.30	0.12	0.02	0.0	1

pattern of the area. High rainfall tends to leach out the basic cations from the soil and could also be as a result of geological materials. Organic matter (GM) content of the soils of the studied area were generally low according to the rating of Landon (1991). However, Okigwe and Egbema both have 1.69% each more than Ihiagwa with a mean value of 0.97%. The low OM in the study area may be attributable to low decomposition rate of organic matter due to the agricultural activities (Adjia *et al.*, 2008) and could also be as a result of their varying parent materials especially, the coastal plain sands (Ihiagwa soils) having a low inherent organic matter content (Obi *et al.*, 2012).

The total nitrogen percentage of all the soils of the area were Low and similar with the mean values of 0.08% for (Okigwe and Egbema) and 0.18% for Ihiagwa soils. The low total nitrogen is typical of the highly weathered soils of the sub-humid tropics, attributed to high nitrogen losses resulting from high rainfall that causes leaching of soil nitrates as well as low organic matter content (Eshett *et al.*, 1989; Brady and Weil, 1999; Opukiri *et al.*, 1991). The low exchangeable bases Mg, K, Na) in the studied soils could be attributed to high rainfall.

Their mean values were similar with Okigwe having 3.9 Cmol/kg, Egbema with 3.23 Cmol/kg and Ihiagwa 3.32 Cmol/kg. According to FAO (2006) these values were low. The result showed no significant difference in the Total Exchangeable acidity of the soils of the area. The soils are acidic due to high concentration of H⁺ and Al³⁺ as a result of leaching (Alekseeva *et al.*, 2009; Barthes *et al.*, 2008).

The effective Cation Exchange Capacity of the soils were low but showed no significant difference. The mean values were low according to the rating of Esu (1991).

Table 3 – 6 shows the macro-aggregate stability of the studied soils. Okigwe had the highest mean values for both water stable aggregate (WSA>0.3) at 5.58% and dry aggregate (DSA>0.3) at 78.07% for Tables 3 and 5, respectively. Also Okigwe recorded the highest Mean Weight Diameter (MWDw) of 0.31% and (MWDa) of 2.01% for Tables 5 and 6 respectively. Egbema recorded 3.71% on WSA>0.3 in table 3 and 48.4% DSA>0.3 with 0.37% and 1.07% MWD in Tables 5 and 6 respectively.

Ihiagwa recorded the least values of 1.77% WSA>0.3 and 14.63% DSA>0.3 in Tables 3 and 4 respectively with 0.44% and 0.16% MWD in Tables 5 and 6.

Generally the soils recorded higher values of DSA>0.3 than in WSA>0.3 and this may be attributed to the low organic matter content of the soils due to leaching and high mineralization rate (Igwe and Nwokocha 2004). The mean Weight Diameter (dry) were smaller with percentage values between 0.25 – 0.45 (Table 5). The difference between these two MWD indices explained the

potential rate of structural deformation of the aggregate (Zobeck *et al.*, 2003).

These low values of MWDw could lead to quick dispersion of the soil during rainfall leading to severe soil erosion (Igwe and Nwokocha, 2004). Although Igwe *et al.*, (1995), Igwe (2003) demonstrated that MWD does not predict well the potential of the soil to erode yet, it is known that finer aggregates flow easily when submerged by water. The potential structural deformation index (PSDI) for the soils ranged from 0.16 to 0.85%. Mbagwu *et al.*, (1991) indicated that the higher the values of PSDI, the more the tendency of the aggregate to disintegrate upon wetting.

Table 7, shows the correlation matrix of some soil physicochemical properties. The WSA>0.3 indicated that it is influenced positively by Mg, TEB and ECEC. It has been stated earlier on that bases such as Ca⁺ and Mg⁺ aid aggregation. Also the higher the amount of TEB, the more acidity can be neutralized in short perspective (Turner and Clark, 1996). This also explained why MWDw correlated negatively with pH. When the MWDw increases it is more stable and reduces erosion. The PSDI is influenced by moisture content. In the tropical climate soils are subjected to frequent wetting and drying cycles, in the short term during the rainy seasons. In terms of climate, the aforementioned long term wetting and drying implications for the aggregate stability of the soils (Igwe and Obalum, 2013).

Also, the characteristics rainstorms and the associated heavy raindrop especially in the humid tropics can have considerable splash effect (Le Bissonais, 2006), Okigwe, Egbema and Ihiagwa as the case study may be and therefore are a force to reckon with in soil aggregate destabilization. It is thus clear that the studied soils are structurally fragile and susceptible to catastrophic erosion (Amezketta, 1999; Bajracharya *et al.*, 1992).

Conclusion

The result of the study generally revealed that the soils are structurally fragile and susceptible to erosion. However, soils of Okigwe and Egbema appeared to be more stable than those of Ihiagwa. The studied soils were sandy especially in Ihiagwa having a mean value of 909.6g kg⁻¹ and least in Okigwe with 629g kg. Soils were generally acidic, ranging from a pH of 3.83 to 5.12. The soils recorded low organic matter content ranging from 0.97 to 1.69% as a result of the parent materials. There was smaller percentage value between the MWDw and MWDd, indicating the potential rate of structures deformation of the aggregate. This leading to severe erosion. The soils of the area should be supported with enough organic manures to increase soil aggregation especially in the soils of Ihiagwa. This will help to improve the water holding capacity, infiltration rates and nutrients retention so as to enhance plant growth and optimum yield.

References

- Adjia, R; Fezeu, W.M.L., Tchatchueng, J.B., Sorho, S. Echevarria, G., Ngassoum, Akamigbo, F.O.R. (1999). Influence of Land use on Soil properties of the humid tropical agro-ecology of Southeastern Nigeria. *Agric. J.* 30: 59 – 76.
- Akamigbo, F.O.R. (2001). Soil Survey, Classification and land use of wetland soils in Nigeria. Proceedings of the 27th Annual Conference of Soil Science Society of Nigeria, Nov. 5 – 9. University of Calabar, Nigeria.
- Alekseeva, T.V, Sokolowska, Z., Hajnos, M., Alekseeva, A.O., Kalinin P.I. (2009). Water Stability of aggregates in subtropical and tropical soils (Georgia and China) and its relationship with the mineralogy and chemical properties. *Eurasian Soil Science* 2009:42:415 – 425.
- Amezketta, E. (1999). Soil aggregate stability: a review. *Journal of sustainable Agriculture* 1483-151
- Attah, L.E. (2010). Physicochemical characteristics of rhizosphere Soils of some cereal crops Amaho Woreda. West Shoa, Ethiopia *Maejo International Journal of Science and Technology* 4(0 1): 93 - 100.
- Brady, N.C and P.R. Wea I., (1990). The nature and property of soils 12th ed. PresenticeHall Inc. New Jersey.
- Bajracharya, R.M, Elliot, W.J, Lal, R (1992) Intern11 erodibt1Y of Ohio soil based on field rainfall simulation. *Soil Science Society of America Journal* 1992; 56:267-272.
- Barthes, B.G., Kouakoua E, Larre-Larriouy, M., Razafimbelo, T.M., de Luca, E.F., Azontonde, A., Neves C.S.V.J., de Freitas P.L., Feller C.L. (2008). Texture and sesquioxide Effects of Water stable Aggregates and Organic matter in some Tropical soils. *Geoderma* 2008; 143:14-25.
- Baye, M.R., Morgan J. and Scholten P.A (2004). Price dispersion in small and in large: evidence from an internet price comparison site. *Journal of Industrial Economics* 52, 463-496.
- Blatt. Harvey and Robert J. T. (1996). *Pedology: Igneous, Sedimentary and Metamorphic*, 2nded, Freeman, pp. 281 — 292.
- Boix-fayos, C., Calvo-Cases, A., Imeson, A.C., Soriano-Soto, M.D. (2001). Influence of Soil Properties on the aggregation of some Mediterranean Soils and the use of aggregate size and stability as land degradation indicators. *Catena* 44(1): 47 – 67.
- Brady, N.C. and Weil R.R. (1999). *The Nature and Properties of Soils*. 14th ed. Prentice Hall: ISBM – 10: 013227938X.
- Bremner, J.M. (1996). Nitrogen-total, In: Sparks, D.I. (ed). *Methods of Soil Analysis part 3. Chemical method*, 2nd Edition. SSSA. Book Series No. 5, SSSA. Madison W.I. 1085 – 1121.
- Calero, N. Barron V, Rorrent, J. (2008). Water dispersible clay in calcareous soils of southwestern Spain *Catena* 2008 74: 22 –30.
- Cambardella, C.A. and Elliot E.T. (1993). Carbon and nitrogen distribution in aggregates from cultivated and native grassland soils. *Soil Sci. Soc. Am. J.* 57: 2071 – 1076.
- Cater E. (1993). Tourism and Natural Resources Management - *Forest Science* 14(2):85 – 90.
- Chenu.CS., I3issonna'S, L. and Arrouays.D. (2000). Organic Matter influence in Characterization of two benchmark soils of contrasting parent materials in Abia State, Southeastern Nigeria. *Global I Pure Applied Sd.*, 16: 23 – 29.
- Cruvinel, P.E., Flocchini, R.G., Crestana, S., Morales, J.R., Miranda, J., Kusko, B.H. and Nielsen, D.R. (1993). Studying some influence of the Aggregate Sizes on some Elements of an Oxisol with PIXE. *Soil Sci.*, 155: 100 – 104.
- Dikinya, O. Hinz, C. and Aylmore G. (2006). Dispersion and re-disposition of fine particles and their effects on saturated hydraulic conductivity. *Australian Journal of Soil Research* 44: 47 – 56.
- Enwezor, W.O., A.C. Ohiri., E.E. Opuwaribo and E. J. Udo (1990). A review of fertilizer use on crops in Southeastern Zone of Nigeria. In: *Literature review on soil fertility investigation in Nigeria*. Fed. Min. of Agric. And Nat. Res. Lagos, Nigeria 2: 49 – 100.
- Eshett, E.L., Otnueto, J.A. and Juo, A.S.R. (1989). Soil Chemical Properties and Mineralogy in relation to Southeastern Nigeria. *J. Agric. Sd.* 112: 377 – 386.
- Esu, I.E. (2005). Characterization, Classification and Management problems of major soils order in Nigeria 27 Inaugural lecture, Nigeria Institute for Agricultural Research, Ahumadu Bellow University, Zaria.
- Esu, L.E. (1991) Detailed Soil Survey of NIHORT Farm at Bunkure, Kano State.
- FAO (Food and Agricultural Organization) (2006), *FAO Production Year book*. Occ, G.W. and Or., (2002). Particle size analysis In: J.H. Dane and G.C Topp (eds). *Methods of soil analysis, part 4. Physical methods of soil survey society of American Hook series*. No: 5 ASA and SSA Madison, W.I., pp. 201-228
- Gee, G.W. and Or, D. (2002). Particle size analysis. In: *method of analysis Dan. D. J. and G. C. Topps (Eds) part 4. Physical method. Soil*

- Science of America book series, No. 5, ASA and SSSA Madison, W.I.P.P. 225 – 295.
- Grossman, R.B. and T.G. Reinesch (2002). Bulk density and Linear extensibility, In: *Methods of Soil Analysis* Dame, J.H. and G.C. Topp (eds), Part 4. Physical methods. Soil Sci Soc. Am. Book Series No. 5ASA and SSSA Madison, WJ. pp. 201 – 228.
- Igwe C.A. (2003) Erodibility Soil of the upper rainforest zone, southeastern Nigeria. *Land Degradation and Development*, 14; 323 – 334.
- Igwe C.A. and Nkwocha D. (2004). Influence of Soil Properties on the Aggregate Stability of a highly Degraded Tropical Soil in Eastern Nigerian mt. *Agro Physics*, 2005, 19; 131 – 139.
- Igwe C.A., Akamigbo F.O.R., and Mbagwu J.S.C (1995). The Use of some Soil Aggregate Indices to assess Potential Soil loss in Soil of southeastern Nigeria. *Tnt. Agro Physics*, 9, 95 100.
- Igwe, C.A. and Obalum, S.E (2013). Microaggregate Stability of Tropical Soils and its Roles in Soil Erosion Hazard Prediction; pp 175 – 187.
- Keinper W.D. and Rosenau R.C. (1986). Aggregate Stability and size distribution. Part 1. In: 'Methods of soil analysis: Physical and Mineralogical Method', 2nd ed: Black C.A. American Society of Agronomy, WI, pp. 425, 432 – 433.
- Landon J.R. (1991). *Booker Tropical Soil manual: A handbook for Soil Survey and Agricultural Land Evaluation in the Tropics and Subtropics*. John Wiley and Sons, New York, ISBN-13: 978-0582005570, PP 474, 849.
- Le Bissonnais, Y. (2006) Aggregate breakdown mechanisms and erodibility. In Lal R (Ed), *Encyclopedia of Soil Science* (2d ed.), Taylor and Francis 2006; 40 – 44.
- M.B (2008). Long term effect of Municipal Solid Waste amendment on Soil heavy metal content of sites used for peri urban agriculture in Ngaoundere Cameroon, *African Journal of Environmental Science and Technology*. Vol. 2 (12) pp. 412 421.
- Mbagwu, J.S.C, Piccolo A., and Spallacip, (1991). Effects of Field Applications of Organic Wastes from different sources on Chemical, Rheological and Structural Properties of some Italian surface soils. *Biore source Tech.*, .7 – 38.
- Mclean. E.O. (1982). Aluminum In: *Methods of Soil Analysis* (ed) C.A. Black Agron No. 9 part 2 Amer. Soc. of Agron. Madison, WJ. 987 – 908.
- NCRS (Natural Resources Conservation Services) (1996). *The United States Department of Agriculture (USDA). Soil Quality Indicators: Aggregate Stability*. Ppl, 18 – 20.
- Nelson, D.W. and Sommers, L.E. (1982). Total Carbon and Organic Matter In: page A.L(Ed). *Society of Agronomy, Madison Wisconsin* Pp. 539 – 579.
- Noellemeyer, E., Frank, F., Alvarez, C., Morazzo, G. and Quiroga, A. (2008). Carbon Contents and Aggregation Related to Soil Physical and Biological Properties under a land-use Sequence in the Semiarid Region of Central Argentina. *Soil Till. Res.*, 99: 179 – 190.
- Obi, C.I., Esu, I.E., Onweremadu, EU., Nwaisi, Felix; Agim Leonard and Chris Emenyonu: (2012). "Characterization and Classification of five Soil Pecloris located within the vicinity of Uyo capital territory, Nigeria. (Unpublished research Article). www.chineduobiwordpress.com.
- Ofomata, G.E.K. (1975). Land Form Regions in Ofomata G.F.K (ed). *Nigeria in map Eastern States*. Ethiope Publishing house, Bcniu City pp. 33 – 34.
- Ogban.P.I., Ekanem, T.P. and Etim, E.A. (2001). Effect of mulching methods on soil properties and growth and yield of maize in south-eastern Nigeria. *Tropical Agriculture(Trinidad)* 78(2): 82 – 89.
- Olson, SR. and L.E. Somniers (1990). Phosphorus. In: Page A.L. (ed) *method of soil analysis part 2 Agron. Monogr, No.9 Madison. W.I.* 403 – 131.
- Onweremadu, E.U., Akamigbo, F.O.R. and Igwe, C.A. (2007). Physical Shrinkage Relationship in Soils of Dissimilar Lithologies in Central southern Nigeria. *Journal of Applied Sciences*, 7(1): 2495 – 2499.
- Opara, C.C. (2009). Soil Microaggregates Stability under Different Land use Types in Southeastern Nigeria. *Catena* 79: 103 – 112.
- Opukiri, S.B., Zuafo K. and Duoglas D.C. (1991). The influence of Urea fertilizer on yield of upland rice (*Oryza sativa*) on acid soils in Rivers State, Nigeria. *Journal Crop, Soil Forestry*. Vol. 1, pp 42 – 43.
- Orajaka.S.O. (1975). Geology, In: Ofomata GJ.K. (ed) *Nigeria in Map Eastern Slate*. Ethiope Publishing house, Benin City. Pp 5 – 7.
- Osuji, G.E. and Onweremadu, E.U., (2005). Aggregation of some Mediterrean Soil and the use of Aggregate Size.
- Six, J., Elliot, E. T. and Paustian, K. (2000). Soil macroaggregate turnover and microaggregate formation: A mechanism for C sequestration under no-tillage agriculture. *Soil Biol. Biochem.* 32: 2099 - 2103
- Six. J. Bossuyt, H. Degryze S., Deneff, K. (2004). A history of research on the link between

- (micro) aggregate stability in red soils from subtropical China, *Catena*; 81: 240 – 248.
- Thomas, G.W. (1996). Soil pH, soil acidity. In: *Methods of Soil analysis part 3. Chemical method* L.D. Sparks (eds) SSA book Series pp. 159 – 165
- Thomas, G.W. (1982). Exchangeable basis. In: *Methods of analysis, part 2* page A L. Miller RH: Keeney, D.R. (eds), Pp. 159 - 165. American Society of Agronomy Madison Wisconsin.
- Turner, R.C. and Clark J.S. (1996). Lime potential in acid clay and suspensions. *Trans. Comm. IL & IV. Int Soc. Soil Sciences*, pp 20 —215.
- Zobeck, T.M., Popham, T.W., Skidmoe, E.L., Lamb, J.A., R.E., (2003). Aggregate mean diameter and wind erodible soil predictions using dry aggregate size distribution *Soil Science Soc. Amer. J.*, 67, 425 - 436. College of tropical Agriculture and Human Resources (CTHAR) (2007). Soil Nutrient Management for Maui County; high Weathered Tropical (Oxisol). www.ctharhawaii.edu/oc/.SCM20.pdf